

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PATENT SPECIFICATION

(11) 1 330 829

DRAWINGS ATTACHED

- (21) Application No. 23426/72 (22) Filed 18 May 1972
(31) Convention Application No. 144 499 (32) Filed 18 May, 1971 in
(33) United States of America (US)
(44) Complete Specification published 19 Sept. 1973
(51) International Classification A61K 27/12
(52) Index at acceptance
A5B 750 756 757 758 75Y 764



(54) LAYERED BOLUS FOR ANIMAL HUSBANDRY PROVIDING FOR IMMEDIATE AND SUSTAINED RELEASE OF MEDICAMENT

(71) We, SMITH KLINE & FRENCH
LABORATORIES, of 1500 Spring Garden
Street, City of Philadelphia, Commonwealth
of Pennsylvania, 19101, United States of
America, a corporation organized under the
laws of the Commonwealth of Pennsylvania,
one of the United States of America, do hereby
declare the invention, for which we pray
that a patent may be granted to us, and the
method by which it is to be performed, to be
particularly described in and by the following
statement:—

This invention relates to a dosage form to
be employed in veterinary practice, particularly
for the administration of therapeutic substances
to ruminants such as cattle and sheep. More
specifically, this invention relates to a layered
bolus which provides both a rapid and sustained
release of a therapeutically active substance
in a single unit dose.

The sustained release bolus employed for
veterinary practice is known to the patent art.
These dosage forms have sufficient dimensions
as to density and weight to stop in the rumeno-
reticular sac and remain there rather than pass
into the alimentary tract and be eliminated
intact. The physical form of these dosage units
is retained over a long period of time in the
sac while the therapeutically active substance
is slowly released by erosive or solubilization
action within the rumeno-reticular sac. In
other words, regardless of the therapeutically
active substance to be administered, the bolus
itself must comply with the physical requirements
as to density and weight. In order to meet
the proper requirements for density and weight,
iron or any other relatively high density matrix
may be employed in the preparation of the bolus.
Specification No. 866924 discloses a veterinary
bolus which provides for the biologically active
substance over an extended period of time. In
the case of a therapeutic substance such as, for
example, a sulfonamide or an antibiotic, the patent
reveals it would be possible to provide a blood level

of three months or more. The above noted
patent also discloses that the rate of release is
accomplished by varying the firing temperature
used in the preparation of the pellets. Extremely
high temperatures are employed to heat pellets
intended to remain in the rumeno-reticular sac
of animals for a prolonged period of time. The
prolonged medication provided by the process of
Specification No. 866924 can last from weeks to
in excess of a year and is desired in order to
prevent retardation of growth and development
of the animal, i.e., more of a prophylactic type
of medication than curative.

Instead of having the medication remain for
months or a year as taught by the above noted
patent, it is often necessary to dose a sick
animal several weeks before slaughtering with
a sulfonamide or other such medicament. When
dealing with food producing ruminants, any
medication given in therapeutic doses should
be essentially clear of the tissues before the
animals are slaughtered. United States Patent
No. 3,507,952 discloses an improved sustained
release bolus for veterinary use which meets
the above release requirement. The patent
reveals a bolus having a predictable and
controlled sustained release pattern of the
medicament over a relatively short period, i.e.,
the bolus disintegrates in the rumen in about
ten days and the animal tissues are clear of the
medicament in three weeks.

However, there is still one major problem
which these prior art boluses fail to solve. It
takes at least twelve hours post administration
for any of the known sustained release sulfon-
amide boluses to show therapeutic plasma
levels. The recommended minimal therapeutic
sulfonamide level is 5 mg./100 ml. when ad-
ministering these long acting sulfonamides to
ruminants, additional intravenous or regular
sulfonamide tablets must also be administered
to provide an initial high level. In brief, con-
current treatment of ruminants with either a
parenteral or tablet dosage form plus the sus-

tained release bolus is necessary to insure rapid therapeutic blood levels as well as prolonged therapy.

5 Scours, which is a highly infectious disease in calves the first few days after birth, is a typical example where immediate antibiotic medication is necessary. This disease is caused by microorganisms and due to the acuteness of the illness, it is imperative that medication
10 be made available to the animal from the dosage form as soon as possible. The calf has need for both an immediate medication to eliminate the bacterial invasion and a sustained release effect to treat the diarrhea symptomatically.

15 It is therefore the object of the present invention to provide a single dosage form which provides for both a rapid therapeutic blood level and prolonged therapy which is essential in antibiotic treatment of ruminants. This dosage form would eliminate the need of a concomitant parenteral dose.

It is a further object of this invention to provide a single dosage form for food producing ruminants which increases the rapidity of obtaining adequate blood levels and decreases
25 post treatment withdrawal time required with prolonged action sulfonamides.

Now in accordance with this invention there is provided a sustained release veterinary
30 layered bolus comprising an immediate release layer and, contiguous therewith, a sustained release layer, said immediate release layer having a disintegration time in the rumen of less than thirty minutes and comprising sufficient of a medicament rapidly to
35 provide a therapeutic level of said medicament in the blood of the ruminant animal to which the bolus is administered and said sustained release layer comprising a filler matrix of such a density that the bolus will lodge in the rumeno-reticular sac of the ruminant animal and sufficient of said medicament that, at the
40 rate of disintegration of said sustained release layer, a therapeutic level of said medicament is maintained in the blood of the ruminant animal to which the bolus is administered for at least 72 hours from the time of dosage.

The layered bolus of this invention provides
45 for plasma levels higher than the recommended minimal therapeutic level (5 mg./%) in less than half the time it requires the prior art bolus to reach this level. The bolus is further advantageous in that it can be administered to food producing ruminants for therapeutic effect several weeks before slaughtering and the medicament present will be essentially clear of the animal tissues at the time of
50 slaughtering, i.e., post treatment withdrawal time is decreased.

60 The bolus of this invention therefore solves the prior art disadvantage of not being able to obtain both an immediate and sustained release of a medicament in a single dose and

fill a commercial need of such a dosage unit in the veterinary field.

Layered tablets for human administration are well known to the pharmaceutical art. However, there is no prior art disclosing a layered bolus of the magnitude and having both immediate and sustained release characteristics noted above as necessary for certain
65 70 veterinary diseases.

The differences between the prior art layered tablet and sustained release layered bolus of this invention are clearly apparent from the ensuing description taken together with the accompanying drawing wherein:

Figure 1 is a side elevation of the prior art layered tablet;

Figure 2 is a side elevation of a layered
75 80 bolus of the present invention;

Figure 3 is a top plan view of the layered bolus;

Figure 4 is an end view of the layered bolus shown in Figure 2;

Figure 5 is a transverse sectional view as seen along the line 5—5 of Figure 3.

Referring specifically to Figure 1, layered tablet 2 has an immediate release layer 4 and a sustained release layer 6. This tablet represents one of the largest prior art layered tablets. The tablet weighs approximately 1.0 grams, has a diameter of 0.503 inches and a thickness of 0.22 inches.

In Figures 2 to 5 the sustained release layered bolus of this invention is disclosed. Bolus 10 weighs 25.97 grams and has an immediate release layer 12 and a sustained release layer 14. The length of the bolus taken horizontally along Figure 2 is approximately 2.733 inches. The width of the bolus as measured vertically in Figure 3 is 0.864 inches and the thickness of the bolus as measured vertically in Figure 4 is 0.630 inches.

A further view of immediate release layer 12 and sustained release layer 14 is presented in the cross section as represented in Figure 5.

In brief, the average layered bolus as embraced by this invention and disclosed in Figure 2 weighs 26 times more than the closest prior art layered tablets. Other dimensions such as thickness and length range from 3 to 6 times greater than presently available layered tablets. In order to prepare the layered bolus of this invention, a specially modified tablet machine is necessary.

The method for the preparation of a bolus according to the present invention comprises preparing separate immediate and sustained release granulations containing a medicament and a filler which at least in the sustained release granulation is of such a density that the bolus will lodge in the rumeno-reticular sac of the ruminant animal, placing said granulations in separate hoppers and selectively compressing said granulations into a layered bolus.
115 120 125

The sustained release effect is accomplished by employing in the sustained release layer of our bolus a lubricant in certain critical percentage ranges as detailed in United States Patent No. 3,507,952.

Tablet disintegrants such as, for example, starch may be used to facilitate disintegration of this layer.

The lubricant employed in the sustained release layer can be any of the more common tablet water insoluble lubricants, such as, for example, magnesium stearate, sodium stearate, calcium stearate, powdered stearic acid, talc, paraffin, cocoa butter, graphite, lycopodium or combinations thereof. The preferred lubricants are fatty acid derivatives especially the stearates, such as, for example, magnesium stearate, sodium stearate, calcium stearate and stearic acid. In order to achieve a definite predictable sustained release pattern, the lubricant material should be present from about 1% to about 3.5% by weight of the sustained release layer. Most advantageously, the lubricant will be present from 2.5% to 3.5% by weight of the sustained release layer.

Exemplary of the dense fillers which may be employed as a matrix are iron powder, calcium sulfate dihydrate, Portland cement, plaster of Paris, magnesium oxyiodide cement or mixtures thereof. The filler will be present from 5% to 95% by weight of the total solids. Preferably the filler will be present from 25% to 75%.

When binders or granulating agents are necessary to insure adequate cohesiveness of the bolus, natural gums and gum constituents such as, for example, acacia, tragacanth, agar, and pectin may be employed. Further, exemplary of binders would be, for example, cellulose esters, polyvinyl-pyrrolidone and proteinaceous material such as, for example, gelatin, casein and zein.

It will be evident to one skilled in the tableting art that the binder is a standard pharmaceutical tool commonly used and is not an essential aspect of this invention, therefore the amount of binder can be varied.

The veterinary layered bolus in accordance with this invention and a method for its preparation will be further illustrated by the following specific example:

EXAMPLE

A granulation is prepared from the following powders:

Sustained Release Granules		
Ingredients	Gms./Bolus	
Sulfamethazine, U.S.P.	13.000	
Calcium Sulfate, Dihydrate	7.000	
Polyvinylpyrrolidone	1.125	
F.D. & C. Yellow #5 Dye	0.005	
F.D. & C. Blue #1 Dye	0.005	
Purified Water	Variable	
Magnesium Stearate	0.213	

The polyvinylpyrrolidone, yellow and blue dyes are dissolved in water. The sulfamethazine and calcium sulfate, dihydrate powders are mixed and granulated with the polyvinylpyrrolidone dye solution. The mixture is screened and dried overnight at 120°F. The dried granulation is passed through a No. 8 U.S. Standard screen and the magnesium stearate is added to the granulation with mixing.

Immediate Release Granules

Ingredients	Gms./Bolus	
Sulfamethazine, U.S.P.	3.0000	
Calcium Carbonate, Precipitated, U.S.P.	0.0860	
Glycine, N.F.	0.4290	
Starch, U.S.P.	0.4290	
Polyvinylpyrrolidone (PVP)	0.2150	
F.D. & C. Yellow #5 Dye	0.0009	
Purified water, U.S.P.	variable	
Starch, U.S.P.	0.2574	
Alginic Acid	0.2574	
Diocetyl Sodium Sulfosuccinate, N.F.	0.0129	

The sulfamethazine, calcium carbonate, glycine and starch powders are thoroughly mixed. The mixture is granulated with an aqueous solution of the PVP and yellow dye. The wet mixture is passed through a No. 4 U.S. Standard screen on an oscillating granulator onto drying trays. The screened mixture is then dried at 120°F. for 12 hours. The dried granulation is passed through a No. 8 U.S. Standard screen on the Stokes oscillating granulator and the remainder of the starch is added together with alginic acid and dioctyl sodium sulfosuccinate.

The immediate and sustained release granulations are placed in separate hoppers in a modified Stokes DD2—11HH tableting machine using oblong punches having a size of 0.860" x 2.725" and concaved on a 1-3/4" diameter wheel, 0.110" deep with a 0.10" flat edge. The sustained release granules are first fed into the die with the lower punch adjusted at such a depth to give the desired fill weight. The upper punch then tamps the sustained release granules just sufficiently to form a cohesive layer. The die then moves to the second hopper containing the immediate release granules. These granules are fed into the die on top of the sustained release layer wherein both granulations are then fully compressed to the desired hardness.

The final product is a compressed oblong layered bolus comprising a mixture of a therapeutically active substance, an inert dense filler and from 1.0% to 3.5% of a lubricant. The bolus provides for a controlled release pattern providing both an immediate and sustained release of the medicament. The finished bolus will weigh from 10.0 grams to 50.0 grams and have a density up to 8.0. Advantageously the finished bolus will weight from 20.0 grams to

- 40.0 grams and have a density of 1.5 to 5.0. The length of the bolus will be from 1.5 inches to 3.0 inches. The immediate release layer completely disintegrates within a half hour and rapidly provides a therapeutic blood level of the therapeutically active substance. The sustained release layer disintegrates within 48 hours and provides a controlled sustained release of the medicament maintaining a therapeutic level for at least 72 hours. The medicament is essentially clear of the tissues in approximately two weeks.
- The layered boluses of this invention containing sulfamethazine were tested *in vivo* to determine the plasma levels of free sulfonamide in cattle and also to observe drug residues in tissues over a period of time. The experimental subjects for this study were twelve Hereford heifers. Each animal was dosed as close as possible to 220 mg./kg. of body weight. Blood was drawn aseptically from the jugular vein in heparinized vacutainers at 0, 6, 12, 24, 48, 72, 96, 120, 144, 168, 192, 216, 240, and 264 hours inclusive after dosing and analyzed for sulfonamide.
- Three animals were killed at 48 hours, three at 144 hours, three at 240 hours, and three at 264 hours after administration of boluses and in each case both kidneys and 1.0 kg. of each of the following tissues collected and immediately frozen: skeletal muscle, fat and liver. The frozen tissues were later assayed for the presence of the sulfonamide.
- The results of these *in vivo* tests are presented in the following tables. Table 1 represents plasma concentrations of sulfamethazine after administration of the layered bolus.

TABLE 1

Mg. Sulfamethazine per 100 ml. Plasma

Animal No.	Time (Hours)								
	0	6	12	24	48	72	96	120	144
254	0.00	7.53	10.57	15.58	13.17	6.73	1.72	0.24	0.10
255	0.00	6.02	9.63	12.89	9.80	—	—	—	—
256	0.00	5.49	6.94	6.88	8.44	7.97	4.65	1.50	0.35
258	0.00	6.73	9.38	12.92	11.58	5.95	1.92	0.35	0.11
259	0.00	6.24	8.74	11.73	10.80	3.78	0.72	0.11	0.00
260	0.00	6.14	8.98	9.41	8.44	—	—	—	—
261	0.00	5.66	9.29	14.28	10.87	8.22	2.98	0.36	0.11
263	0.00	5.40	7.56	7.94	7.49	5.42	2.74	0.51	0.11
264	0.00	4.85	7.84	13.21	11.53	4.87	1.05	0.12	0.00
265	0.00	4.83	5.65	6.60	10.50	8.13	2.87	0.51	0.25
267	0.00	5.99	7.71	8.55	5.81	—	—	—	—
268	0.00	7.08	10.66	12.81	7.54	2.35	0.61	0.11	0.00
MEAN	0.00	6.00	8.61	11.07	9.66	5.94	2.14	0.42	0.11

TABLE 2

Free Sulfonamidide Tissue Levels

Time of Tissue Harvest	PPM Free Sulfonamide			
	Fat	Muscle	Kidney	Liver
48 hr.	6.30	17.33	15.70	20.50
144 hr.	0.10	<0.20	0.52	0.62
240 hr.	<0.04	<0.04	<0.04	<0.04
264 hr.	<0.04	<0.04	<0.04	<0.04

The above results indicate that mean levels of 6.0 mg./100 ml. were present in six hours with 5.94 mg./100 ml. still remaining after 72 hours. Both of these levels are above the recommended minimal therapeutic level of 5 mg./100 ml. The animal tissues showed residues of less than 0.04 ppm sulfonamide ten days after administration.

In summary, the above standard animal test procedure demonstrates that when a layered sustained release bolus according to this invention is administered to cattle therapeutic blood levels were rapidly achieved and maintained for days. Further the withdrawal time of the medicament from the tissues has been decreased. The reported tissue residues require a 21 day withdrawal before meat from treated animals may be used for human consumption. As noted above, the sulfamethazine was relatively clear of the tissues ten days post administration.

Although the invention is specifically exemplified using sulfamethazine since the advantages are particularly suited to this medication, it will be appreciated that it is useful with any solid medicament which it is desired to provide in combined immediate and sustained release form. The medicament may be in the form of a base, salt or ester and may be soluble or insoluble in nature. Thus, for example, the medicament may be a sulfonamide derivative such as sulfamethazine, sulfathiazine, a tranquilizer such as chlorpromazine, an antibiotic such as chloramphenicol, tetracycline or penicillin, an anthelmintic such as piperazine phosphate, an anti-bloat agent such as organopolysiloxane, hormone growth supplements such as stilbesterol and many more medicaments such as vitamins and those used in the treatment of bacterial enteritis such as furazolidone.

WHAT WE CLAIM IS:—

1. A sustained release veterinary layered

bolus comprising an immediate release layer and contiguous therewith, a sustained release layer, said immediate release layer having a disintegration time in the rumen of less than thirty minutes and comprising sufficient of a medicament rapidly to provide a therapeutic level of said medicament in the blood of the ruminant animal to which the bolus is administered and said sustained release layer comprising a filler matrix of such a density that the bolus will lodge in the rumeno-reticular sac of the ruminant animal and sufficient of said medicament that, at the rate of disintegration of said sustained release layer, a therapeutic level of said medicament is maintained in the blood of the ruminant animal to which the bolus is administered for at least 72 hours from the time of dosage.

2. A veterinary bolus according to claim 1 wherein the medicament is a sulfonamide or antibiotic and the filler matrix is calcium sulfate dihydrate or iron powder.

3. A veterinary bolus according to claim 2 wherein the medicament is sulfamethazine and the filled matrix is calcium sulphate dihydrate.

4. A veterinary bolus according to claim 1 as hereinbefore described and as illustrated in Figures 2 to 5 of the accompanying drawings.

5. A method of preparing a veterinary layer bolus according to claim 1 having a sustained release layer contiguous with an immediate release layer which comprises preparing separate immediate and sustained release granulations containing a medicament and a filler which at least in the sustained release granulation is of such a density that the bolus will lodge in the rumeno-reticular sac of the ruminant animal, placing said granulations in separate hoppers and selectively compressing said granulations into a layered bolus.

6. A method according to claim 5 wherein the medicament is a sulfonamide or antibiotic and the dense filler is calcium sulphate dihydrate or iron powder.

7. A method according to claim 6 wherein the medicament is sulfamethazine and the filler matrix is calcium sulfate dihydrate. wherein a bolus according to claim 3 is administered orally to the calves. 5
8. A method of treating scours in calves

R. A. A. HURST,
Agent for the Applicants.

Printed for Her Majesty's Stationery Office by the Courier Press, Leamington Spa, 1973.
Published by the Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from
which copies may be obtained.




FIG. 1. (PRIOR ART)




FIG. 2.

(PRESENT INVENTION)




FIG. 3.




FIG. 4.




FIG. 5.